

**A Survey of Values Associated with Riparian
Conditions of a Stream Tributary to the
Green/Colorado River**

**Quentin D. Skinner
Thomas A. Wesche**

**Michael A. Smith
Sue Lowry**

**1989
WWRC-89-48**

In

**Practical Approaches to Riparian Resource
Management: An Educational Workshop**

**Quentin D. Skinner, Michael A. Smith
and Thomas A. Wesche
Department of Range Management
University of Wyoming
Laramie, Wyoming**

**Sue Lowry
Wyoming State Engineer's Office
Cheyenne, Wyoming**

PRACTICAL APPROACHES TO

**RIPARIAN RESOURCE
MANAGEMENT**

AN EDUCATIONAL WORKSHOP

MAY 8-11, 1989
BILLINGS, MONTANA

Edited By

Robert E. Gresswell
Bruce A. Barton
Jeffery L. Kershner

Copy Editor

Kenneth A. Hashagen, Jr.

Printed By

U. S. Bureau of Land Management
222 North 32nd Street
P.O. Box 36800
Billings, Montana 59107

Support for the Workshop Provided By

Montana Chapter of the American Fisheries Society
Montana Riparian Association
Western Division American Fisheries Society
Society for Range Management
U. S. Bureau of Land Management
U. S. Forest Service, Northern Region
U. S. Fish and Wildlife Service
U. S. National Park Service, Yellowstone National Park

A Survey of Values Associated with Riparian Conditions of a Stream Tributary to the Green/Colorado River

QUENTIN D. SKINNER, MICHAEL A. SMITH, AND THOMAS A. WESCHE

*Department of Range Management
University of Wyoming
Box 3354
Laramie, Wyoming 82071 USA*

SUE LOWRY

*Wyoming State Engineer's Office
Cheyenne, Wyoming 82002 USA*

Abstract.—Sediment and salt are important but often ignored contributions to nonpoint pollution in the semi-arid western USA. Accelerated erosion and sediment/salt in stream flow are often thought to be caused by livestock grazing on rangeland in less than excellent condition. Poor condition streams are often downcut and show active bank rilling, bank sloughing, and piping. Lateral movements of gullies away from downcut streams through upland ecosystems are, in addition, contributing sources for increased sediment/salt pollution. Equally important, when downcut streams occur, associated riparian vegetation is often reduced in areal distribution, disappears, or is replaced by pseudoriparian plant species. Loss of riparian vegetation and the hydrologic support of limited riparian ecosystems may further reduce stream channel stability and sediment/salt trapping efficiency of the stream. These nonpoint source pollutants are transported to the stream by overland flow and ground water. With the value of stream and riparian zone stability in mind, the University of Wyoming, U. S. Bureau of Land Management, ranchers, and state agencies initiated a study in the cold desert of south central Wyoming.

This study was located along Muddy Creek between Rawlins and Baggs, Wyoming and was designed to:

(1) Evaluate the use of wire faced dams to enhance the restoration of a degraded stream channel reach and thus the adjoining riparian zones.

(2) Evaluate the amount of sediment and salt stored by riparian zones along Muddy Creek.

(3) Evaluate the surface and ground water regimes associated with degraded, improving, and improved stream channel reaches along Muddy Creek.

(4) Evaluate vegetation response between a degraded and improving stream reach along Muddy Creek.

(5) Compare economic returns of riparian zones supported by different stream channel conditions to the cost of restoring them using sediment, salts, vegetation, and stored ground water.

Preliminary results show that riparian zones filter sediment from streamflow. Diversion and flood irrigation, flooding by beaver dams, and water spreading controlled by dikes have been successful in creating a surface aquifer/riparian zone in a desert ecosystem that is valuable for reducing nonpoint pollution and for prolonged storage of water during late summer months. Sediment can be trapped and vegetation production increased using wire faced dams in desert streams. Because of confining bed material under and along the existing channel, storage of water and creation of surfaced aquifer with dams was limited to the sediment storage capacity within the existing downcut stream channel. Both water diversion and damming of streamflow show improvement in riparian zone vegetation and control of nonpoint pollution when compared to a degraded stream reach.

Our study shows that stored sediment, trapped salts, increase in vegetation, and perhaps stored water in riparian zones, when using wire faced dams, will provide benefits sufficient to cover cost of construction and maintenance for federal and state agencies as well as municipalities but not the private producer. However, water diversion, water spreading, and creating a surface aquifer may increase riparian vegetation and store water sufficient to pay back the investment for these water development practices used in the past by private land owners.