

**Response of Mountain Stream Channels and
Associated Riparian Areas to Flow Regulation**

Thomas A. Wesche

Steven W. Wolff

Quentin D. Skinner

1989

WWRC-89-49

In

**Practical Approaches to Riparian Resource
Management: An Educational Workshop**

**Thomas A. Wesche and Steven W. Wolff
Wyoming Water Research Center
University of Wyoming
Laramie, Wyoming**

**Quentin D. Skinner
Department of Range Management
University of Wyoming
Laramie, Wyoming**

PRACTICAL APPROACHES TO

**RIPARIAN RESOURCE
MANAGEMENT**

AN EDUCATIONAL WORKSHOP

MAY 8-11, 1989
BILLINGS, MONTANA

Edited By

Robert E. Gresswell
Bruce A. Barton
Jeffery L. Kershner

Copy Editor

Kenneth A. Hashagen, Jr.

Printed By

U. S. Bureau of Land Management
222 North 32nd Street
P.O. Box 36800
Billings, Montana 59107

Support for the Workshop Provided By

Montana Chapter of the American Fisheries Society
Montana Riparian Association
Western Division American Fisheries Society
Society for Range Management
U. S. Bureau of Land Management
U. S. Forest Service, Northern Region
U. S. Fish and Wildlife Service
U. S. National Park Service, Yellowstone National Park

Response of Mountain Stream Channels and Associated Riparian Areas to Flow Regulation

THOMAS A. WESCHE AND STEVEN W. WOLFF

*Wyoming Water Research Center
Box 3076, University Station
Laramie, Wyoming 82071 USA*

QUENTIN D. SKINNER

*Department of Range Management
P. O. Box 3354
University of Wyoming 82071 USA*

Abstract.—The need for channel maintenance flow releases below water development projects in the central Rocky Mountain region is currently being debated. The objective of our study was to measure the change in hydraulic geometry of mountain stream channels in response to water diversion. Channel width, depth, width to depth ratio, cross-sectional area, and conveyance capacity were measured above and below diversion structures on 20 stream reaches in Wyoming and Colorado to determine the extent of channel adjustment since regulation. Ocular estimates of riparian vegetation density were also made. Diversion structures ranged in age from 12 to 106 years and reduced streamflow up to 90%.

Our preliminary analysis using paired *t*-tests indicates no significant ($P > 0.05$) differences in channel dimensions above and below diversions on steep (slope $> 4.0\%$) and moderate (slope 1.5 to 4.0%) gradient channels. Low gradient (slope $< 1.5\%$) channels responded to streamflow depletion by significantly reducing their depth, area, and conveyance capacity. In a majority of cases, we observed no differences in vegetation density above and below diversion structures.

Our preliminary results indicate that moderate to high gradient mountain stream channels located in the forest snowpack zone may be maintained with reduced streamflow regimes. Low gradient channels, where encroachment and aggradation appear more likely to occur, should be the primary focus of channel maintenance flow studies.